## Virginia Asthma Action Plan

School Division: Date of Birth Name **Effective Dates** GREEN means Go! Use CONTROL medicine daily Health Care Provider Provider's Phone YELLOW means Caution! Add RESCUE medicine Parent/Guardian Parent/Guardian Phone Parent/Guardian Email: RED means DANGER! Get help from a doctor now! Additional Emergency Contact Contact Phone Contact Email: Asthma Triggers (Things that make your asthma worse) Asthma Severity Last Flu Pneumonia ☐ Colds ☐ Smoke (tobacco, incense) ☐ Pollen ☐ Dust Shot: Shot: □ Intermittent □ Strong odors □ Mold/moisture 1 1 11 ☐ Pests (rodents, cockroaches) ☐ Stress/Emotions ☐ Exercise Persistent: 

Mild 

Moderate □Gastroesophageal reflux □ Season (circle):Fall, Winter, Spring, Summer □ Severe Take these CONTROL (PREVENTION) Medicines EVERY Day ☐ No control medicines required. Always rinse mouth after using your daily inhaled medicine. You have ALL of these: \_ puff (s) MDI with Spacer \_\_\_ times a day Breathing is easy No cough or wheeze \_ nebulizer treatment (s) \_\_\_ times a day Inhaled Corticosteroid Can work and play , take by mouth once daily at bedtime • Can sleep all night Leukotriene antagonist For asthma with exercise, ADD: Peak flow in this area: \_\_\_ puffs with spacer 15 minutes before exercise Fast acting Inhaled β-agonist (More than 80% of Personal Best) For nasal/environmental allergy, ADD: Personal best peak flow: \_\_ spray (s) per nostril \_\_\_\_ times a day \_, use \_\_\_ Nasal corticosteroid Yellow Zone: Caution! — Continue CONTROL Medicines and ADD RESCUE Medicines You have ANY of these: \_ puffs with spacer every \_\_\_\_ hours as needed Inhaled b-agonist • First sign of a cold \_ nebulizer treatment (s) every \_\_\_\_\_ hours as needed · Cough or mild wheeze Inhaled b-agonist Tight chest ☐ Other Problems sleeping, working, or playing Call your Healthcare Provider if you need rescue medicine for more than 24 Peak flow in this area: hours or two times a week, or if your rescue medicine doesn't work to (60%-80% of Personal Best) **DANGER!** — Continue CONTROL & RESCUE Medicines and GET HELP! Red Zone \_, \_\_ puffs with spacer **every 15 minutes**, for **THREE** treatments You have ANY of these: Inhaled 8-agonist • Can't talk, eat, or walk well nebulizer treatment every 15 minutes, for THREE treatments Medicine is not helping Inhaled 8-agonist Call your doctor while administering the treatments. · Breathing hard and fast □ Other Blue lips and fingernails IF YOU CANNOT CONTACT YOUR DOCTOR: Tired or lethargic Ribs show Call 911 for an ambulance, Peak flow in this area: or go directly to the Emergency Department! to (Less than 60% of Personal Best) REQUIRED SIGNATURES: SCHOOL MEDICATION CONSENT AND HEALTH CARE PROVIDER ORDER I give permission for school personnel to follow this plan, administer medication FOR CHILDREN/YOUTH and care for my child and contact my provider if necessary. I assume full responsibility for providing the school with prescribed medication and delivery/ CHECK ALL THAT APPLY: monitoring devices. I approve this Asthma Management Plan for my child. Student has been instructed in the proper use of all of his/her asthma medications, and in my opinion, <u>CAN CARRY AND SELF-ADMINISTER HIS or</u> HER INHALER AT SCHOOL. PARENT/GUARDIAN SCHOOL NURSE/DESIGNEE Student is to notify his/her designated school health officials after using Virginia Asthma Action Plan approved by the Virginia Asthma Coalition (VAC) 4/11 Student needs supervision or assistance to use his/her inhaler. Based on NAEPP Guidelines and modified with permission from the D.C. Asthma Action Plan via District of Columbia Department of Health, DC Control Asthma Now, and District of Columbia Student should NOT carry his/her inhaler while at school.

DATE

MD/NP/PA SIGNATURE: \_

Blank copies of this form may be reproduced or downloaded from www.virginiaaasthma.org